



## Field application of pure polyethylene microplastic has no significant short-term effect on soil biological quality and function

Robert W. Brown<sup>a,\*</sup>, David R. Chadwick<sup>a</sup>, Harriet Thornton<sup>b</sup>, Miles R. Marshall<sup>a</sup>,  
Shuikuan Bei<sup>c</sup>, Marco A. Distaso<sup>a,d</sup>, Rafael Bargiela<sup>a,d</sup>, Karina A. Marsden<sup>a</sup>, Peta L. Clode<sup>e</sup>,  
Daniel V. Murphy<sup>f</sup>, Saskia Pagella<sup>a,b</sup>, Davey L. Jones<sup>a,f</sup>

<sup>a</sup> School of Natural Sciences, Bangor University, Bangor, Gwynedd, LL57 2UW, United Kingdom

<sup>b</sup> Graduate School of the Environment, Centre for Alternative Technology, Machynlleth, Powys, SY20 9AZ, United Kingdom

<sup>c</sup> Centre for Resources, Environment and Food Security, College of Resources and Environmental Sciences, China Agricultural University, Beijing, 100193, China

<sup>d</sup> Centre for Environmental Biotechnology, Bangor University, Bangor, Gwynedd, LL57 2UW, United Kingdom

<sup>e</sup> Centre for Microscopy, Characterisation & Analysis and School of Biological Sciences, The University of Western Australia, Perth, WA, 6009, Australia

<sup>f</sup> SoilsWest, Centre for Sustainable Farming Systems, Food Futures Institute, Murdoch University, Murdoch, WA, 6150, Australia

### ARTICLE INFO

#### Keywords:

Plastic pollution

Metabolomics

Toxicology

Soil quality

Environmental impact

### ABSTRACT

Plastics are now widespread in the natural environment. Due to their size, microplastics (MPs; defined as particles <5 mm) in particular, have the potential to cause damage and harm to organisms and may lead to a potential loss of ecosystem services. Research has demonstrated the significant impact of MPs on aquatic systems; however, little is known about their effects on the terrestrial environment, particularly within agroecosystems, the cornerstone of global food production. Soil biology is highly responsive to environmental perturbation and change. Hereby, we investigated the effect of pure low-density polyethylene (LDPE) MP loading (0, 100, 1000, or 10000 kg ha<sup>-1</sup>) on soil and plant biological health in a field environment over a cropping season. Our results showed that MP loading had no significant effect ( $p > 0.05$ ) on the soil bacterial community diversity (as measured by amplicon sequencing of bacterial 16S rRNA gene), the size and structure of the PLFA-derived soil microbial community, or the abundance and biomass of earthworms. In addition, metabolomic profiling revealed no dose-dependent effect of MP loading on soil biogenic amine concentrations. The growth and yield of wheat plants (*Triticum aestivum* L., cv. Mulika) were also unaffected by MP dose, even at extremely high ( $\geq 1000$  kg ha<sup>-1</sup>) loading levels. Nitrogen (N) cycling gene abundance before and after N fertiliser application on the MP loaded experimental plots showed relatively little change, although further experimentation is suggested, with similar trends evident for soil nitrous oxide (N<sub>2</sub>O) flux. Overall, we illustrate that MPs themselves may not pose a significant problem in the short term (days to months), due to their recalcitrant nature. We also emphasise that most MPs in the environment are not pure or uncontaminated, containing additives (e.g. plasticisers, pigments and stabilisers) that are generally not chemically bound to the plastic polymer and may be prone to leaching into the soil matrix. Understanding the effect of additives on soil biology as well as the longer-term (years to decades) impact of MPs on soil biological and ecological health in the field environment is recommended.

### 1. Introduction

The use of plastics is globally ubiquitous due to their low cost, malleability, and durability; however, inappropriate disposal has led to their progressive accumulation in the environment (Geyer et al., 2017). To date, much of plastic and microplastic (MPs; particles <5 mm in size) pollution research has focused on freshwater and marine systems, where

the negative effects of plastics on organism health and loss of ecosystem function is now becoming well documented (Avio et al., 2017; Sharma and Chatterjee, 2017). However, plastics are also rapidly being identified as a threat to terrestrial ecosystems, yet their potential effects remain largely unexplored (de Souza Machado et al., 2019).

In agroecosystems, plastic entry may occur through a variety of pathways, with the most common including (i) the use, and

\* Corresponding author. School of Natural Sciences, Bangor University, Gwynedd, Bangor, Gwynedd, LL57 2UW, UK.

E-mail address: [rob.brown@bangor.ac.uk](mailto:rob.brown@bangor.ac.uk) (R.W. Brown).

incorporation of plastic mulch films to improve plant growth and reduce moisture loss (Huang et al., 2020; Sun et al., 2020; R. Qi et al., 2020); (ii) the addition of municipally-derived organic fertilisers, digestates or compost (Watteau et al., 2018); (iii) the application of biosolids (van den Berg et al., 2020); (iv) the accumulation of slow-release fertiliser coatings (Katsumi et al., 2021) and (v) atmospheric deposition (Allen et al., 2019) (vi) irrigation from polluted sources (Bläsing and Amelung, 2018). The drive for food security and sustainable intensification has led to an inevitable increase in plastic loading to soils globally. For example, the annual input of plastics into agricultural soils is estimated to be between 63–430 and 44–300  $\times 10^3$  t in Europe and North America, respectively, and potentially exceeding  $1.3 \times 10^6$  t annually for China (Jian et al., 2020; Nizzetto et al., 2016a). Globally, this greatly surpasses the extrapolated annual mass discharge of MPs to ocean surface waters, predicted to be  $9.3 \times 10^7$ – $2.36 \times 10^8$  tonnes (Nizzetto et al., 2016a, 2016b; Sebillle et al., 2015). Primary MPs (MPs manufactured for a specific application, e.g. clothing microfibres; de Falco et al., 2019) may be applied through waste streams (i.e. biosolids application), due to their difficulty of removal (Cole et al., 2011). In contrast, secondary MPs are formed through degradation and disintegration of larger plastic pieces (Cole et al., 2011; Rocha-Santos and Duarte, 2015), such as agricultural mulch films (Piehl et al., 2018). Both primary and secondary MPs are likely to influence the ecology, health and function of soils, potentially having similar negative effects to those extensively reported in marine ecosystems, e.g. organismal ingestion leading to oxidative stress and assimilation of endocrine-disrupting chemicals, and subsequent reduced growth and reproduction, as well as bioaccumulation up the food chain (Galloway and Lewis, 2016; Kim et al., 2017). Although, bioaccumulation is likely to be less of an issue comparatively, due to the relatively smaller size of soil-dwelling fauna.

Soil is an extremely valuable and non-renewable resource and provides range of ecosystem services, not least the provisioning of food resources (Comerford et al., 2013; Kopittke et al., 2019). Maintaining soil health and quality is therefore key for agricultural and anthropogenic sustainability (Hou et al., 2020). Soil quality is often broadly defined as the capacity of a soil to function (Karlen et al., 1997). Traditional measurements of soil quality are based on physical or chemical soil properties, with little exploration of soil biology (Bünermann et al., 2018). However, the fertility and productivity of soil are not simply a function of soil physical and chemical characteristics, and recently a more holistic view has been proposed (Rinot et al., 2019). Soil biology is a crucial mediator and driver of many processes linked to nutrient cycling, plant health, and soil productivity (Lal, 2016). It is highly responsive to changes in management and environmental conditions and is often associated with functional change (Lehman et al., 2015a,b). Research has shown that MPs can have significant negative effects on soil microbial community composition (Guo et al., 2020; Zang et al., 2020; Zhang et al., 2019), enzymatic activities and nutrient cycling (Fei et al., 2020; Huang et al., 2019; Yi et al., 2021), mesofaunal health (Huerta Lwanga et al., 2016; Lahive et al., 2019; Lin et al., 2020), plant health (de Souza Machado et al., 2019; Zang et al., 2020), and greenhouse gas (GHG) emissions (Ren et al., 2020; Sun et al., 2020), all of which will impact the soils ability to function effectively. However, most studies to date have been laboratory or mesocosm based, over relatively short sampling periods (weeks) and in many cases at unrealistic MP doses, which may not accurately reflect processes occurring at the field scale (Fidel et al., 2019).

This field-based study aimed to assess the effect of different quantities (0, 100, 1000, or 10000 kg ha<sup>-1</sup>) of pure MP loading on the health and function of key soil biological quality indicators over a cropping season, using a range of commonly used biological indicators, as well as the novel use of biogenic amine analysis as indicators of metabolism and N cycling in soil. Loading rates were chosen to represent 'existing', 'normal', 'future', and 'extreme' (or 'hotspot') MP loading to soil (Gao et al., 2019; Huang et al., 2020; R. Qi et al., 2020). Pure MP was chosen as much of the current literature does not disentangle the effect of pure

plastic from the plastic additives for example, UV stabilisers (Stenmarck et al., 2017) and pigments (Gičević et al., 2020). This study aims to serve as a "negative" control, supporting future research on these chemicals and helping to exclude confounding effects that could derive from the particulate nature of the plastic particles. We hypothesised that i) MP loading will have negative effects on all measured aspects of soil biological quality, ii) higher MP loading rates will increase the detrimental impact on soil biology, and iii) crop biomass and yields will be negatively affected by MP loading.

## 2. Materials and methods

### 2.1. Experimental setup


The experiment took place at the Henfaes Agricultural Research Station, Abergwynnregyn, North Wales (53°14'N, 4°01'W). The soil is classified as a sandy clay loam textured Eutric Cambisol, overlying a glacial till, with a temperate-oceanic climate. The mean annual rainfall is 1060 mm and the mean annual temperature is 10 °C. The site has no previous history of plastic pollution or application over the last 50 years (Zang et al., 2020). On April 18, 2019, a randomised plot design was established to create 4 independent replicates ( $n = 4$ ) of each treatment. Each plot (1.4  $\times$  2.85 m) was then treated with LDPE microplastic powder (RXP1003 natural; Resinex Ltd., High Wycombe, UK), at a rate of 0, 100, 1000, or 10000 kg ha<sup>-1</sup> by thorough manual mixing with the top 10 cm of soil. This equated to loading rates of 0%, ~0.1%, ~1%, and ~10% (w/w) (soil bulk density = 1040 kg m<sup>-3</sup>;  $n = 4$ ). The microplastic powder was confirmed to have a very low level of contamination through total carbon (C) and nitrogen (N) analysis using a TruSpec® Analyzer (Leco Corp., Michigan, USA) (Total C, 82.88%  $\pm$  0.03%; Total N, 0.03  $\pm$  0.01%;  $n = 5$ ). LDPE was chosen due to its extensive use in agricultural films (Espí et al., 2006; Rong et al., 2021). Plots were subsequently sown with spring wheat (*Triticum aestivum* L., cv. Mulika) at a rate of 400 plants m<sup>-2</sup>. In line with the fertiliser recommendations for wheat, and taking account of the soil's Soil Nitrogen Supply (SNS) (AHDB, 2018), 120 kg N ha<sup>-1</sup> yr<sup>-1</sup> was applied to the field as NH<sub>4</sub>NO<sub>3</sub> over two applications, 40 kg N ha<sup>-1</sup> on 3rd June and 80 kg N ha<sup>-1</sup> on 3rd July (reflecting the late sowing of the crop). For scanning electron microscopy (SEM), LDPE powder was mounted on adhesive tape, coated with gold, and imaged at 10 kV (Tescan Vega3 SEM). These SEM images illustrate the heterogeneous nature of the MP mixture, both in terms of particle size and surface texture (Fig. 1).

### 2.2. Soil sampling and analysis

The soil was sampled one, two, and six months following MP addition. On each sampling occasion, multiple fresh soil cores per plot ( $n = 12$ ;  $\phi = 1$  cm; depth = 0–10 cm) were randomly sampled and homogenised by hand to obtain a representative plot soil sample. Soil pH and electrical conductivity (EC) were measured on 1:2.5 (w/v) soil-to-distilled water suspensions by submerging standard electrodes. Within 24 h of soil collection, 1:5 (w/v) soil-to-0.5 M K<sub>2</sub>SO<sub>4</sub> extracts were performed. The colorimetric methods of Miranda et al. (2001) and Mulvaney (1996) were used to determine the nitrate (NO<sub>3</sub>-N) and ammonium (NH<sub>4</sub>-N) concentrations in the K<sub>2</sub>SO<sub>4</sub> extracts, respectively. Bulk density cores (0–5 cm, 100 cm<sup>3</sup>) were also collected oven-dried (105 °C, 24 h) before being weighed. Soil characteristics are summarised in Table 1. Climatic data from an adjacent weather station for the sampling period and a timeline of sampling are summarised in Fig. S1.

### 2.3. Phospholipid fatty acid (PLFA) profiling of the microbial community

Soil sampling for PLFA analysis was performed after 2 and 6 months of MP addition. Fresh homogenised soil samples, collected as described in section 2.2, were subsampled for PLFA analysis. The subsampled soil



**Fig. 1.** Scanning electron micrographs of microplastic particles before incorporation into the soil. The images were taken across a range of magnifications (A – 20 µm; B – 50 µm; C – 100 µm; D – 200 µm; E – 200 µm; F – 500 µm). Images illustrate the heterogeneous nature of particle size and surface texture within the powder sample.

**Table 1**

Influence of microplastic (MP) dose and time since application on soil properties. The soil was sampled one, two or six months post microplastic application. Results are expressed on mean dry soil weight basis  $\pm$ SEM ( $n = 4$ ). Letters denote significant differences between treatments ( $p < 0.05$ ).

	1 month post-MP application				2 months post MP application				6 months post MP application			
	0	100	1000	10000	0	100	1000	10000	0	100	1000	10000
MP loading rate (kg ha <sup>-1</sup> )												
pH	6.26 $\pm$ 0.04 <sup>a</sup>	6.23 $\pm$ 0.19 <sup>a</sup>	6.26 $\pm$ 0.14 <sup>a</sup>	6.23 $\pm$ 0.10 <sup>a</sup>	6.49 $\pm$ 0.04 <sup>a</sup>	6.34 $\pm$ 0.15 <sup>a</sup>	6.41 $\pm$ 0.12 <sup>a</sup>	6.47 $\pm$ 0.08 <sup>a</sup>	6.27 $\pm$ 0.11 <sup>a</sup>	6.16 $\pm$ 0.26 <sup>a</sup>	6.14 $\pm$ 0.11 <sup>a</sup>	6.09 $\pm$ 0.08 <sup>a</sup>
EC ( $\mu$ S cm <sup>-1</sup> )	129 $\pm$ 38 <sup>a</sup>	91 $\pm$ 13 <sup>a</sup>	123 $\pm$ 24 <sup>a</sup>	96 $\pm$ 22 <sup>a</sup>	37 $\pm$ 1.9 <sup>a</sup>	36 $\pm$ 2.6 <sup>a</sup>	31 $\pm$ 2.3 <sup>a</sup>	31 $\pm$ 3.5 <sup>a</sup>	55 $\pm$ 2.4 <sup>a</sup>	77 $\pm$ 25 <sup>a</sup>	55 $\pm$ 3.9 <sup>a</sup>	51 $\pm$ 2.6 <sup>a</sup>
NO <sub>3</sub> <sup>-</sup> (mg N kg <sup>-1</sup> )	67.4 $\pm$ 21.7 <sup>a</sup>	18.6 $\pm$ 4.6 <sup>a</sup>	33.4 $\pm$ 14.5 <sup>a</sup>	38.3 $\pm$ 0.70 <sup>a</sup>	5.04 $\pm$ 2.60 <sup>a</sup>	4.96 $\pm$ 3.02 <sup>a</sup>	1.86 $\pm$ 0.09 <sup>a</sup>	1.61 $\pm$ 0.14 <sup>a</sup>	10.4 $\pm$ 4.30 <sup>a</sup>	21.9 $\pm$ 9.32 <sup>a</sup>	15.5 $\pm$ 4.1 <sup>a</sup>	10.2 $\pm$ 1.08 <sup>a</sup>
NH <sub>4</sub> <sup>+</sup> (mg N kg <sup>-1</sup> )	57.5 $\pm$ 16.7 <sup>a</sup>	11.0 $\pm$ 5 <sup>a</sup>	22.1 $\pm$ 10.9 <sup>a</sup>	45.8 $\pm$ 1.6 <sup>a</sup>	1.01 $\pm$ 0.06 <sup>a</sup>	1.11 $\pm$ 0.11 <sup>a</sup>	1.13 $\pm$ 0.05 <sup>a</sup>	0.89 $\pm$ 0.06 <sup>a</sup>	2.64 $\pm$ 0.30 <sup>a</sup>	5.36 $\pm$ 2.09 <sup>a</sup>	3.28 $\pm$ 0.88 <sup>a</sup>	3.00 $\pm$ 1.05 <sup>a</sup>
Bulk density (kg m <sup>-3</sup> )					1014 $\pm$ 11 <sup>a</sup>	1065 $\pm$ 27 <sup>a</sup>	984 $\pm$ 30 <sup>a</sup>	977 $\pm$ 31 <sup>a</sup>	1065 $\pm$ 22 <sup>a</sup>	1106 $\pm$ 48 <sup>a</sup>	1092 $\pm$ 44 <sup>a</sup>	1062 $\pm$ 61 <sup>a</sup>
Bacterial/Fungal PLFA ratio					0.11 $\pm$ 0.01 <sup>ab</sup>	0.11 $\pm$ 0.01 <sup>ab</sup>	0.11 $\pm$ 0.01 <sup>ab</sup>	0.14 $\pm$ 0.02 <sup>a</sup>	0.09 $\pm$ 0.00 <sup>b</sup>	0.10 $\pm$ 0.00 <sup>ab</sup>	0.11 $\pm$ 0.01 <sup>ab</sup>	0.10 $\pm$ 0.01 <sup>ab</sup>
Microbial PLFA biomass ( $\mu$ mol PLFA kg <sup>-1</sup> )					174 $\pm$ 11 <sup>ab</sup>	175 $\pm$ 9 <sup>ab</sup>	162 $\pm$ 3 <sup>a</sup>	190 $\pm$ 16 <sup>ab</sup>	199 $\pm$ 6 <sup>ab</sup>	201 $\pm$ 8 <sup>ab</sup>	197 $\pm$ 6 <sup>ab</sup>	218 $\pm$ 12 <sup>b</sup>
Earthworm biomass (g m <sup>-2</sup> )									92 $\pm$ 9 <sup>a</sup>	54 $\pm$ 6 <sup>a</sup>	71 $\pm$ 24 <sup>a</sup>	79 $\pm$ 22 <sup>a</sup>
Earthworm abundance (individuals m <sup>-2</sup> )									26 $\pm$ 5 <sup>a</sup>	13 $\pm$ 2 <sup>a</sup>	24 $\pm$ 13 <sup>a</sup>	20 $\pm$ 6 <sup>a</sup>

EC – electrical conductivity

was subsequently stored at  $-80$  °C to prevent lipid turnover. Lyophilisation was performed using a Modulyo Freeze Dryer (Thermo Electron Corporation, Waltham, MA, USA) attached to a rotary vane pump (Edwards Ltd., Crawley, UK). Samples were shipped on dry ice ( $-78.5$  °C) to Microbial ID Inc. (Newark, DE, USA) for analysis. The method of [Buyer and Sasser \(2012\)](#) was used for extraction, fractionation and transesterification of samples. Analysis was performed on a 6890 gas chromatograph (GC) (Agilent Technologies, Wilmington, DE, USA) equipped with an autosampler, split-splitless inlet, and flame ionization detector. The system was controlled with MIS Sherlock® (MIDI, Inc., Newark, DE, USA) and Agilent ChemStation software. GC-FID specification, analysis parameters and standards are as described in [Buyer and Sasser \(2012\)](#).

#### 2.4. Biogenic amine extraction and analysis

Biogenic amine extraction was performed 6 months after microplastic addition. Biogenic amines are a subset of the metabolome, key in the processing and cycling of N, which has previously been shown to be

sensitive to changes in biological quality ([Brown et al., 2021](#); [Withers et al., 2020](#)). On this sampling occasion, additional multiple soil cores ( $n = 5$ ;  $\phi = 1$  cm; depth = 0–10 cm) were taken across each plot and homogenised by hand to obtain a representative soil sample. After collection, samples were transferred ( $<1$  h) to a  $-80$  °C freezer to quench metabolic amine turnover. Samples were stored and lyophilised as described in section 2.3. Post-lyophilisation, roots and other debris (e.g. plant litter) were removed and the samples were then ground using a stainless-steel ball mill (MM200, Retsch GmbH, Haan, Germany), to aid in the recovery of biogenic amines. The mill was sterilised between samples by rinsing with deionised water followed by a 70% ethanol solution. Ground soil was transferred to sterile polypropylene 1.5 ml microfuge tubes and shipped, on dry ice ( $-78.5$  °C), to the West Coast Metabolomics Center (UC Davis Genome Center, Davis, CA, USA) for untargeted biogenic amine analysis using hydrophilic interaction chromatography electrospray quadrupole time of flight tandem mass spectrometry (HILIC-ESI QTOF MS/MS).

Briefly, extraction consisted of vortexing ( $\sim 15$  s) a 0.4:1 (w/v) soil-to-3:3:2 (v/v/v) MeCN/IPA/H<sub>2</sub>O solution, before shaking for 5 min at

4 °C, centrifuging (2 min, 14000 g) and recovering an aliquot of the supernatant for analysis. LC/QTOFMS analysis of extracted aliquots was performed on an Agilent 1290 Infinity LC system (G4220A binary pump, G4226A autosampler, and G1316C Column Thermostat) coupled to a SCIEX Triple TOF mass spectrometer, total runtime was 16.8 min. Polar compounds are separated on an Acquity UPLC BEH Amide Column, 13 nm (pore size), 1.7 µm (particle size), 2.1 mm × 150 mm maintained at 45 °C at a flowrate of 0.4 ml min<sup>-1</sup>. Solvent pre-heating (Agilent G1316) was used. The mobile phases consist of: (A) Water, 10 mM ammonium formate, 0.125% formic acid and (b) acetonitrile: water (95/5, v/v), 10 mM ammonium formate, 0.125% formic acid. The gradient was: 0 min 100% (B), 0–2 min 100% (B), 2–7 min 70% (B), 7.7–9 min 40% (B), 9.5–10.25 min 30% (B), 10.25–12.75 min 100% (B), 16.75 min 100% (B).

A sample volume of 1 µl for positive mode and 3 µl for negative mode was used for the injection. Sample temperature was maintained at 4 °C in the autosampler. The mass spectrometer was operated with gas temperatures set to 300 °C and pressure to 345 kPa (curtain gas (CUR) – 2.4 bar; IonSpray Voltage Floating (ISFV) – 4500 V; declustering potential (DP) – 10 V; capillary electrophoresis (CE) – 100V). Electrospray ionization (ESI) performed full scans in the mass range *m/z* 50–1200. The number of cycles in MS1 was 1667 with a cycle time of 500 ms and an accumulation time of 475 ms. Data were collected in both positive and negative ion mode and analysed using MS DIAL, open software for metabolome analysis, as described in Tsugawa et al. (2015). Final curated results were reported as peak heights, internal standards were included, however, these were for quality control and peak correction purposes. Data presented are therefore qualitative and compounds are tentatively identified, as is routine for untargeted analysis (Gertsman and Barshop, 2018). A full compound list is presented in supplementary information with standardised reference nomenclature being generated using RefMet (Fahy and Subramaniam, 2020).

## 2.5. Soil N<sub>2</sub>O flux

A mobile, automated GHG monitoring system, utilising a GC-Electron Capture Detector (8610C, SRI Instruments, CA, USA), as previously described in Marsden et al. (2018), was used to monitor nitrous oxide (N<sub>2</sub>O) fluxes from three of the four replicates for each treatment. Stainless steel chamber bases (50 × 50 cm; 0.25 m<sup>2</sup>) were installed into plots two weeks before MP application, to which chambers (0.0625 m<sup>3</sup>) were tightly secured. A foam strip on the base of each chamber ensured a tight seal. Briefly, the automated sampling system provided eight greenhouse gas flux measurements per 24 h period, per chamber during uninterrupted measurement. Emissions were monitored for 6 months from installation. However, this manuscript focuses on the 2-week periods following initial MP loading, to test whether the background emissions from the soil were perturbed by MP incorporation and the two subsequent N fertiliser application events, respectively, as these periods were likely to produce the greatest fluxes (Bell et al., 2015; Cardenas et al., 2019).

## 2.6. High-throughput sequencing and quantitative PCR analysis

### 2.6.1. 16S rRNA gene sequencing

Soil samples for 16S rRNA gene sequencing were collected after 6 months of MP incorporation. Five soil cores (*n* = 5;  $\phi$  = 1 cm; depth = 0–10 cm) were taken from each plot and homogenised by hand to obtain a representative sample. After collection, samples were passed through a 2 mm sieve and subsequently transferred (<1 h) to a –80 °C freezer for pre-extraction storage. Genomic DNA was extracted by mechanical lysis from 0.25 g soil per sample using a DNA Soil Fecal/Soil Microbiome Kit (ZymoResearch, CA, USA). Quality and concentration of extracted DNA were assessed by agarose gel electrophoresis (AGE) using a Qubit 4.0 Fluorometer dsDNA BR Assay Kit (Life Technologies, United States). Libraries of 16S rRNA gene amplicons were created using primers for

rRNA marker genes (identical to those described in Distaso et al. (2020)), specifically for the V4 region of the 16S rDNA targeting bacteria and archaea (515F/806R), were prepared as previously described in Fadrosch et al. (2014). PCR was performed using a ViiA7 qPCR system (Applied Biosystems, MA USA). Thermocycling conditions were: initial denaturation at 95 °C for 3 min, followed by 25 cycles of 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 30 s with a final elongation at 72 °C for 5 min. Purified amplicons were then quantified using the aforementioned Qubit 4.0 Fluorometer, pooled in equimolar amounts and the final pool was run on the Illumina MiSeq platform (Illumina Inc., CA).

### 2.6.2. Bioinformatic analysis

The previously described protocols of Fadrosch et al. (2014) and Distaso et al. (2020) were used to process raw sequencing reads. In total, 214,318 raw sequencing reads were produced. Briefly, data pre-processing extracted the barcodes from sequences, and subsequently cleaned primer sequences using tagcleaner. Barcodes and sequences were then re-matched using in-house python scripts and the resulting filtered reads analysed using QIIME v1.9.1. Erroneous sequences and Chimeras were removed using quality filtering during demultiplexing, and ChimeraSlayer, respectively, both were implemented in QIIME. The libraries were demultiplexed based on the different barcodes. Sequences were then classified into operational taxonomic units (OTUs) combining *de novo* and reference-based methods (open-reference OTU generation algorithm) using the SILVA reference database version 132 (Yilmaz et al., 2014). Here, OTUs were determined using an open-reference OTU picking process, where reads are clustered against a reference sequence collection and any reads which do not hit the reference sequence collection are subsequently clustered *de novo*, only OTUs with a minimum coverage of 20 were included in the analysis. Chloroplast and Mitochondrial reads were removed from the OTU count. Sequencing read files analysed in this study can be accessed from the National Center for Biotechnology Information (project PRJNA762001).

### 2.6.3. Quantitative PCR of N cycling functional genes

Samples for quantitative PCR (qPCR) of N cycling functional genes were collected on the 3rd July (pre-N fertiliser application) and on the 15th July (12 days post-N fertiliser application). On each occasion five soil cores (*n* = 5;  $\phi$  = 1 cm; depth = 0–10 cm) were taken per plot and homogenised by hand to obtain a representative sample. After collection, samples were passed through a 2 mm sieve and subsequently transferred (<1 h) to a –80 °C freezer for pre-extraction storage. Samples were extracted for NO<sub>3</sub>-N and NH<sub>4</sub>-N, as described in section 2.2. DNA was extracted by mechanical lysis from 0.25 g soil per sample using a DNEASY Powersoil kit (Qiagen, Hilden, Germany). The quality and concentration of extracted DNA were assessed by AGE.

To obtain the standard curves for qPCR assays, functional genes (urease (*ureC*), archaeal ammonia oxidation (AOA-*amoA*), bacterial ammonia oxidation (AOB-*amoA*), complete nitrification (*comammox*), nitrite reductase (*nirK*; *nirS*), nitrous oxide reductase (*nosZ*) and nitrogenase iron protein (*nifH*)) were amplified using the primers listed in Table S1 qPCR was performed using a QuantStudio 7 System (Applied Biosystems, Waltham, United States). The thermocycling conditions are for each gene are summarised in Table S1. For each gene, a high amplification efficiency of 92–105% was obtained, the R<sup>2</sup> values were >0.99 and no signal was observed in the negative controls. The copy numbers for each sample of soil DNA were calculated based on comparison with the standard curve. qPCR was performed using a QuantStudio 7 System (Applied Biosystems, Waltham, United States). Results were subsequently normalised by the extracted DNA concentration for each sample to account for differences in extraction efficiencies within samples and raw results are included in supplementary information.

## 2.7. Earthworm abundance and biomass

Earthworm abundance and weight were assessed after 6 months.

Briefly, a 0.018 m<sup>3</sup> (0.3 × 0.3 × 0.2 m) pit was dug in a randomly selected location in each experimental plot. Soil from the pit was placed into a tray and thoroughly manually sorted, and earthworms collected. All earthworms were counted (abundance) and weighed (biomass). Abundance is expressed as individuals m<sup>-2</sup> and biomass as fresh weight biomass m<sup>-2</sup>.

## 2.8. Wheat harvest data

Spring wheat was harvested at full maturity, 5 months after sowing. The harvest protocol consisted of hand cutting, with shears, a 1 × 2.85 m strip, through the center of each experimental plot, to remove edge effects. Samples were then dried (85 °C, 48 h). For each harvested sample, ears were removed from stems and each were weighed. Ear and stem weight were subsequently added to calculate a total wheat biomass dry weight per plot or biomass yield. Biomass yield was used as it is highly related to grain yield and gives an overall indicator of plant health (Damisch and Wiberg, 1991). After drying, harvested wheat seeds were separated, weighed and ground, and subsequently analysed for total C and N using a TruSpec® Analyzer (Leco Corp., St. Joseph, MI, USA) and a C:N ratio calculated.

## 2.9. Statistical analysis

All statistical analysis was run using R v 4.0.3 (R Core Team, 2021) unless otherwise stated. With all graphical analysis being constructed in 'ggplot2' (Wickham, 2016) unless otherwise stated. A significance level of  $p < 0.05$  was used for all analyses.

Normality and homogeneity of variance of the chemical and physical soil properties of the treated Eutric Cambisol were assessed using Shapiro-Wilk's test and Levene's test, respectively. For data that did not conform to parametric assumptions even after using log<sub>10</sub> transformation (NO<sub>3</sub>-N, NH<sub>4</sub>-N, EC and PLFA Fungal:Bacterial ratio) a Kruskal-Wallis test (stats package; R Core Team, 2021) was used to assess the similarities between MP treatments and sampling dates, otherwise a one-way ANOVA (Analysis of variance) was used (for pH, bulk density and total PLFA biomass). The results for this are summarised in Table 1. A one-way ANOVA was also used to assess treatment variations in wheat biomass data (total aboveground biomass, stem and leaf biomass, ear biomass and harvested wheat seed C:N ratio) and earthworm data (abundance and biomass).

The 'vegan' (Oksanen et al., 2020) and 'ggplot2' (Wickham, 2016) packages were used to construct NMDS (Non-metric multidimensional scaling) analysis of the PLFA community based on Bray-Curtis dissimilarities. All PLFAs detected were used in the analysis, to represent the whole microbial community. This was followed by computation of an ANOSIM (Analysis of similarities) to identify differences in dispersion between centroids of groups as determined by MP loading rate, or time of sampling. Fungal-bacterial ratios and Gram positive to Gram negative ratios were calculated by summing the FA biomarkers for the respective groups (summarised in Table S2). Total biomass was calculated by summing the concentration of PLFAs recovered.

Fluxes of N<sub>2</sub>O for each chamber were calculated using the methods described in Scheer et al. (2014). The linear slope of N<sub>2</sub>O concentrations over time included either three or four data points. N<sub>2</sub>O fluxes for each two-week period (post-MP and fertiliser application, respectively) were graphically analysed. Trapezoidal integration was used to calculate cumulative N<sub>2</sub>O emissions for each treatment, these were tested for significance using for Kruskal-Wallis tests, after failing parametric assumptions.

Bacterial observed OTU richness was tested for significant differences using ANOVA. The evenness of the 16S community was also calculated using Pielou's evenness (Jost, 2010) and tested for significant differences using ANOVA. NDMS, followed by an ANOSIM (Analysis of similarities) was used to test statistically whether there was a significant difference between groups of sampling units between treatments

(β-diversity).

N cycling gene abundance, before and after a N fertilisation event was analysed using mixed effect models with the 'lme4' package (Bates et al., 2015). We considered MP loading rate and sampling time and their interaction as fixed effects and individual plots as temporal random effects. For each variable, residuals from each model were tested for normality, autocorrelation and heteroscedasticity using graphical tools. For all genes, a log<sub>10</sub> conversion was found to improve the fitness of all models. An ANOVA was then run on each model to test treatment effects, significant results were further explored using a Tukey adjusted post-hoc test using the 'emmeans' package (Lenth, 2021). Pre- and post-fertilisation soil NO<sub>3</sub>-N and NH<sub>4</sub>-N concentrations were analysed by ANOVA.

MetaboAnalyst v5.0 (Chong et al., 2018; Pang et al., 2020) was used for the analysis of biogenic amine data. First, normalisation was performed using generalised logarithm transformation (glog) and Pareto scaling. Normalised data was subsequently used for heatmap creation (using Euclidean distance and Ward clustering algorithms). One-way ANOVA was also performed to identify significant differences in compound concentrations between treatments.

Also, we acknowledge that, being a field trial, a high level of representative replication (i.e., replication with large enough plot sizes) is difficult to obtain, which could potentially impact the statistical power of the study. However, on calculating the statistical power of the parametric statistics used here all were ≥0.99, with the expectation of bacterial OTU evenness (power = 0.05), thus this result should be interpreted with caution.


## 3. Results

### 3.1. 16S bacterial community

In total, 7179 bacterial operational taxonomic units (OTUs), were identified across all 16S rRNA gene reads. There was little variation in the proportional abundance of OTUs between the different MP treatments with Proteobacteria (Gram-negative) and Actinobacteria (Gram-positive) being the most abundant phyla (Fig. 2A). There were no significant differences between bacterial OTU richness ( $F_{(3,12)} = 0.32$ ,  $p > 0.8$ ) (Fig. 2B) or evenness ( $F_{(3,12)} = 1.74$ ,  $p > 0.2$ ) (Fig. 2C) across the different treatments, as tested by ANOVA. Equally, the NMDS ordination shows no clear separation or divergence in soil bacterial communities between the MP treatments and the unamended control (Fig. 2D). Lastly, we found no significant differences in bacterial β-diversity between the treatments, as confirmed by ANOSIM analysis ( $p > 0.8$ ).

### 3.2. PLFA-derived community

The fungal-bacterial ratio of PLFAs remained similar across all treatments, there was a significant difference between the 2 months post-application 10000 kg ha<sup>-1</sup> and the 6 months post-application 0 kg ha<sup>-1</sup> MP loading rates, with the latter having a higher prevalence of bacteria (Table 1). Total PLFA biomass was also similar across all treatments, with a significant difference between the 2 months post-application 1000 kg ha<sup>-1</sup> and the 6 months post-application 10000 kg ha<sup>-1</sup> MP loading rates, the latter having a higher PLFA biomass yield. NMDS analysis was used to show the clustering of all soil-derived PLFA compounds, under MP treatments, 2 and 6 months after initial MP application (Fig. 3). Overall, the different MP treatments separated by sampling date, with a clear separation between the 2 and 6-month points. The PLFA derived community was also more closely grouped at the 6-month sampling point. Results of the PERMANOVA confirmed that there was no significant difference in group dispersion between MP loading treatments ( $p > 0.2$ ). There was, however, a significant difference in group dispersion between sampling times ( $p < 0.001$ ), additionally there was no interaction effect between MP loading and sampling time ( $p > 0.9$ ).



**Fig. 2.** 16S rRNA gene sequenced bacterial community in response to different microplastic doses ( $n = 4$ ). A) Proportionate abundances of major phyla within each microplastic loading rate. B) Boxplot of observed bacterial OTU richness against microplastic loading rate ( $n = 4$ ). C) Boxplot of bacterial OTU evenness against microplastic loading rate ( $n = 4$ ). D) Non-metric multidimensional scaling (NMDS) ordination plot of bacterial OTU community composition across microplastic loading rates.

### 3.3. *N* cycling genes

The presence and abundances of eight genes involved in the N cycle, specifically *ureC*, *amoA* (AOA, AOB, and comammox), *nirK*, *nirS*, *nosZ* and *nifH*, (functions are summarised in Fig. S2), were assayed by qPCR before and after an N fertilisation event. We acknowledge that the primers used to amplify the functional genes (e.g. *ureC*) do not target all of the community. In most cases, gene abundance was not greatly affected by either MP loading rate or sampling time (i.e. pre- and post-N fertilisation) (Fig. 4, Table S3). However, ANOVA showed that there were significant differences for *nirK* ( $F_{(3,12)} = 4.6$ ,  $p < 0.05$ ) and *nosZ* ( $F_{(3,24)} = 3.2$ ,  $p < 0.05$ ) abundance, respectively, by MP loading. For both *nirK* and *nosZ* gene abundance, LMS post-hoc analysis showed a significant difference between 100 kg ha<sup>-1</sup> and 1000 kg ha<sup>-1</sup> MP loading ( $p < 0.05$ ). For AOB, ANOVA also showed a significant interaction effect between MP loading rate and sampling time ( $F_{(3,24)} = 3.5$ ,  $p < 0.05$ ). LMS post-hoc analysis showed that there were significant differences between 0 kg ha<sup>-1</sup> and 1000 kg ha<sup>-1</sup> MP loading, pre fertilisation ( $p < 0.05$ ) and between 0 kg ha<sup>-1</sup> MP loading, pre fertilisation, and 10000 kg ha<sup>-1</sup> MP loading post fertilisation ( $p < 0.05$ ). Concentrations of soil NO<sub>3</sub>-N ( $F_{(1,12)} = 16.6$ ,  $p < 0.01$ ) and NH<sub>4</sub>-N ( $F_{(1,12)} =$

22.0,  $p < 0.01$ ) were significantly higher post-fertilisation (Fig. 4E and F).

### 3.4. N<sub>2</sub>O flux

Kruskal-Wallis analysis showed that there were no significant differences between cumulative N<sub>2</sub>O fluxes for the 2 week period following initial MP application ( $H_{(3)} = 0.74$ ,  $p = 0.9$ ), or the first ( $H_{(3)} = 4.6$ ,  $p = 0.2$ ) and second fertiliser ( $H_{(3)} = 3.6$ ,  $p = 0.3$ ) application events. Fluxes over each period are summarised in Fig. 5.

### 3.5. Biogenic amines

Untargeted biogenic amine analysis identified a total of 112 tentatively identified compounds. Of these known compounds detected, none showed statistically significant differences between treatments. There were no clear grouping or responses within the biogenic amine data (Fig. 6). The samples were characterised by a wide range of compounds (Fig. S3) but predominated by amino acids and peptides.




Fig. 3. NMDS plot of the PLFA profile for each microplastic soil treatment. Ellipses represent 95% confidence intervals for each treatment.

### 3.6. Soil properties including inorganic N

Overall, there were no significant differences in soil chemical properties (pH, EC,  $\text{NO}_3\text{-N}$  and  $\text{NH}_4\text{-N}$ ) associated with the MP treatment as tested by ANOVA or Kruskal Wallis ( $p > 0.1$ ). Trends in the data show some natural variation in all soil properties measured throughout the season (summarised in Table 1).

### 3.7. Earthworms abundance and biomass

Earthworm abundance and biomass were not significantly affected by MP loading. All earthworms identified in the samples were endogenic. Overall, there were no significant differences between total earthworm biomass ( $F_{(3,12)} = 0.63, p > 0.6$ ) or earthworm abundance ( $F_{(3,12)} = 0.85, p > 0.4$ ; Table 1).

### 3.8. Plant biomass

Plant biomass was not significantly affected by MP loading, however, yields of this field trial were lower than the typical wheat yields for the year (DEFRA, 2019). There were no significant differences between total above ground plant biomass ( $F_{(3,12)} = 0.09, p > 0.9$ ), stem and leaf biomass ( $F_{(3,12)} = 0.08, p > 0.9$ ), ear biomass ( $F_{(3,12)} = 0.09, p > 0.9$ ), or harvested seed C:N ratio ( $F_{(3,11)} = 0.03, p > 0.9$ ; Fig. 7).

## 4. Discussion


### 4.1. 16S bacterial community response to MP addition

Soil microorganisms are vital to soil functioning and are considered the most sensitive indicator of soil quality, due to their ability to rapidly respond to changing environmental conditions (Bünemann et al., 2018; Lau and Lennon, 2012; Schimel, 2018). Therefore, despite a significant amount of functional redundancy (Jia and Whalen, 2020), substantial shifts in the microbial community are likely to represent a change in soil function (Lehman et al., 2015a,b). This study showed that after 6 months of pure microplastic addition to previously uncontaminated soil, there was no significant change in the proportional abundance of the bacterial community (Fig. 2A), bacterial richness (Fig. 2B), evenness, or bacterial community compositional divergence ( $\beta$ -diversity) (Fig. 2D). To contextualise this, a previous study at the same site, showed significant changes in the microbial community under biochar application over similar time scales (Jones et al., 2012).

Currently, the effect of MPs loading on soil microorganisms is unclear. Our findings are contradictory to several studies with loading rates equating to  $\leq 5\%$  (lower than the highest loading rate here of 10%), which observed significant effects of microplastic (e.g. LDPE; Huang et al., 2019), polyvinyl chloride (PVC; Yan et al., 2020), and combined PE and PVC (Fei et al., 2020; Seeley et al., 2020) addition on the soil bacterial community, particularly richness, evenness, and diversity. However, H. Y. Chen et al. (2020) and Judy et al. (2019) showed various microplastic additions had no significant effects on the microbial community over short time periods (70 d and a loading rate of 2% and 9 months and a loading rate of up to 10%, respectively). Additionally, Ren et al. (2020) reported mixed but largely positive effects of MP (at a loading rate of 5%) on the microbial community (increase in richness and diversity) in a fertilised soil over a 30 d period, although the microorganisms may have reacted to the fertiliser addition and not the MPs. Based on these studies it is evident that the type of plastic incorporated into the soil will dictate the biological and ecological effects exhibited, therefore a further study of the effect of different types of plastic, and combinations of plastics are required to fully understand any impact on soil health.

### 4.2. Effect of MP loading on soil PLFAs

PLFAs give a representation of the living soil microbial biomass and provide a snapshot of soil community structure and abundance at the time of sampling. NMDS clustering of PLFA microbial community shows a large amount of overlap between MP loading rates implying community structure had not changed significantly (Fig. 3). This is contrary to previous microcosm studies that have shown significant shifts in PLFA derived microbial community even under relatively low levels (from 1%) of MP loading (Zang et al., 2020). MPs are a recalcitrant C pool and are only likely to become bioavailable as a viable C source over long time periods (years to decades) with the aid of natural abiotic degradation (hydrolysis, photo-oxidation or thermal oxidation) (Ángeles-López et al., 2017; Chamas et al., 2020) and to a lesser extent biological degradation (e.g. earthworms) (Huerta Lwanga et al., 2016). This biochemical inertness in the short to medium term is unlikely to cause major shifts in microbial communities. In terms of soil physical properties, MPs have been suggested as a new and distinct microbial habitat, for example for biofilm colonisation and formation (McCormick et al., 2014; Zhang et al., 2019), potentially leading to a change in the microbial community. However, this was not observed in this study as there was no significant community divergence in MP treatments from



**Fig. 4.** N cycling gene soil abundances pre- and post-N fertiliser application ( $n = 4$ ). A) Urease-associated gene *UreC*, B) Free N fixation associated gene *nifH*, C) Nitrification-associated genes, the *amoA* gene of; i) AOA, ii) AOB, iii) *comammox*, D) Denitrification-associated genes; i) *nirK*, ii) *nirS*, iii) *nosZ*, E) Soil nitrate, F) Soil ammonium. All genes abundances were normalised by extracted DNA quantities to account for differences in microbial biomass and transformed by  $\log_{10}$ . Soil nitrate and ammonium are reported by dry soil weight ( $n = 4$ ).







Fig. 5.  $\text{N}_2\text{O}$  fluxes from soil upon; A) initial MP loading, B) N fertilisation event one ( $40 \text{ kg N ha}^{-1}$  equivalent), C) N fertilisation event two ( $80 \text{ kg N ha}^{-1}$ ), by MP loading treatment. In each panel, the line represents the mean flux ( $n = 3$ ) and the shaded area represents the upper and lower bounds of the SEM.

control plots in either 16S bacterial community or PLFA derived microbial community. The SEM (Fig. 1) illustrates that the MP powder used here is not porous or cavity-containing and therefore may not offer an attractive habitat for microbial colonisation (Or et al., 2006). Additionally, we would dispute this theory, as studies with biochar, a similarly recalcitrant C source, have shown that microbial colonisation is very sparse, concluding that even after several years biochar did not provide a substantial habitat for soil microbes (Quilliam et al., 2013). However, this requires confirmation with experimental evidence for

MPs.

Separation between all MP loading treatments groups between the two sampling points (2 months and 6 months post MP addition) illustrated a distinct temporal shift in the structure of the microbial community. Seasonal as well as cropping associated shifts in the PLFA composition in soil have been observed (Duncan et al., 2016; Ferrari et al., 2015; Moore-Kucera and Dick, 2008). These shifts are generally associated with membrane adaptation to changing environmental stress levels (for example, temperature, moisture or nutrient availability),



**Fig. 6.** Influence of microplastic application rate on the biogenic amine (BA) concentration in soil. Heatmap showing expression profiles of soil treatments based on the top 50 most significant know BAs identified by ANOVA ( $p < 0.03$ ). BAs are clustered using Euclidean distance and Ward linkage. Data was normalised using a  $\log_{10}$  transformation and Pareto scaling. The colour of samples ranges from red to blue, indicating metabolite concentration z-score; numbers 3 to  $-3$  on the scale bar indicate the number of standard deviations from the mean. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

resulting in physiological community change (Blagodatskaya and Kuzakov, 2013; Bossio and Scow, 1998). It is likely the observed change in the soil PLFA community between sampling points may be due to natural seasonal changes (for example the difference in soil moisture, illustrated in Fig. S1).

#### 4.3. Effect of N cycling gene abundance pre- and post- N fertilisation

Within agroecosystems, N availability is often considered the predominant limiting factor in plant growth (Vitousek and Howarth, 1991) and the second most limiting factor after C in microbial growth (Kuyper et al., 2018; Buchkowski et al., 2015). Microbial uptake, assimilation, and cycling of mineral and organic N is key to soil function, and as such N cycling processes (mineralisation, nitrification, and denitrification) have been used as sensitive and ecologically relevant indicators of soil quality and ecological stability (Bünemann et al., 2018; Iqbal et al., 2020). Changes in the abundance of the key regulatory functional genes involved in these processes are likely to indicate changes in soil function. However, there is little evidence of how MPs could affect soil N cycling (Iqbal et al., 2020). Overall, this study showed little change in the abundance of N cycling functional genes between pre- and post-inorganic N addition under all MP loading rates. Genes that did

differ significantly in abundances between treatments were denitrification associated (*nirK* and *nosZ*) and nitrification associated (*AOB amoA*). For both denitrification associated genes, lower abundances were displayed within the 1000 kg ha<sup>-1</sup> treatment compared to the 100 kg ha<sup>-1</sup> treatment (Fig. 4C), with no effects on abundances at either higher or lower MP loading rates. *AOB amoA* gene abundance was significantly lower than control levels in the 100 kg ha<sup>-1</sup> treatment pre-fertilisation and 10000 kg ha<sup>-1</sup> treatment post-fertilisation. The general trend in N cycling gene abundances showed variability pre-fertilisation. Post-fertilisation this variability was reduced and gene abundances were more even across all MP loading treatments, while soil inorganic N was significantly increased post-fertilisation (Fig. 4).

N fertilisation has been shown to have a mixed effect on N cycling genes (Tosi et al., 2020). Effects are highly dependent on the nature of the N source applied (inorganic or organic), with inorganic sources of N having a much weaker effect than organic sources of N, as well as the fertiliser duration, crop rotation, and pH (Ouyang et al., 2018). The results of this study show that there were no large changes in soil N cycling functional genes in the presence of MP loading. Although there may have been several further factors influencing N gene abundance, for example when fertiliser was applied the soil was very dry (Fig. S1), preventing soil biology from accessing the additional N. Equally, as




Fig. 7. Effect of microplastic application rate on above-ground wheat biomass ( $n = 4$ ). A) Total above-ground biomass, B) Stem and leaf biomass, C) Ear biomass and D) Seed C:N ratio.

alluded to above, C is the primary limiting factor for soil microbiology, if the community was already C limited then it is unlikely that there would be significant growth or change stimulated by N addition. Studies have shown that MPs have the potential to affect N cycling processes, for example repression of key N cycling enzymes (e.g. leucine-aminopeptidase and N-acetyl- $\beta$ -glucosaminidase (Awet et al., 2018; Bandopadhyay et al., 2020)) and N hydrolysis (Huang et al., 2019). However, N cycling is a key soil function, particularly in agricultural soil, and the longer-term impacts of MPs on should be explored in more detail.

#### 4.4. Effect of MP loading on soil $N_2O$ flux

$N_2O$  is a potent greenhouse gas, with a global warming potential (GWP) 298 times larger than carbon dioxide ( $CO_2$ ) and it is a stratospheric ozone-depleting substance (Stocker, 2014). In soil, it is primarily produced by the biological pathways of nitrification and denitrification. As such it can be used as a functional indicator of soil biological quality at an ecosystem processes scale (Bünemann et al., 2018). Therefore, understanding whether MP addition influences soil  $N_2O$  fluxes will be key to understanding their overall environmental impact. It has been shown that MPs may reduce soil  $N_2O$  emissions by inhibiting the microbial phyla associated with N cycling genes (Ren et al., 2020; Rillig et al., 2021), although results vary depending on the type of MP applied and environmental conditions (Shen et al., 2020; Sun et al., 2020).

While chambers in this study included plant and soil, the plant contribution of  $N_2O$  is minimal (Chang et al., 1998), therefore we focussed on the soil contribution. Here,  $N_2O$  flux from the soil after MP and fertiliser applications, respectively, were very low (Fig. 5).  $N_2O$  fluxes are commonly observed after fertiliser application (up to  $250 \mu g N_2O-N m^{-2} h^{-1}$ ; Carswell et al., 2018), however, we observed none. Equally, there were no differences between fluxes between MP loading levels (Table S4). However, it is difficult to attribute this low flux directly to the microplastic application, particularly as control plots also exhibited small fluxes. Notably, much of the sampling period was dry (Fig. S1), this is likely to have suppressed  $N_2O$  emission, as water filled pore space (WFPS) was too low to allow the development of the anaerobic ‘hotspots’ required for  $N_2O$  production (via denitrification) and emission (Barrat et al., 2020; Dobbie and Smith, 2001). We therefore recommend further field-based measurement of MPs effect on  $N_2O$

and other GHGs (particularly  $CO_2$  and methane ( $CH_4$ )), under a range of climatic conditions and soil types.

#### 4.5. Biogenic amines as effected by MP loading

BAs are low molecular weight organic bases synthesised by prokaryotes and eukaryotes in the soil, mainly through decarboxylation of amino acids or amination and transamination of aldehydes and ketones. In a food context, BAs are often seen as undesirable due to their potentially toxic properties (Mah et al., 2019), in this sense they are potential food quality indicators (Ruiz-Capillas and Herrero, 2019). However, there is also evidence that BAs have a role in quorum sensing in the gut between bacteria and host organisms (Hughes and Sperandio, 2008; Sudo, 2019).

There has been little exploration of BAs in the soil system specifically. But it is generally understood that increased N availability in the soil will increase the number of BAs synthesised both by soil biota and plants (Pérez-Álvarez et al., 2017). Equally, homospermidine biosynthesis has been proposed as a stress regulator in rhizobia (Fujihara, 2009). In this study, one of the first to profile the soil BAs, we found no significant change in the BA amine profile of soil applied with MPs compared to control values, 6 months after initial MP application (Fig. 5, Fig. S3). A large range of compounds were extracted, many of which have putative functions including 5'-methylthioadenosine, an inhibitory by-product of methionine metabolism, which can be processed to salvage biogenically available sulphur (North et al., 2017). As well as abscisic acid, a plant hormone that regulates many aspects of plant growth, including development, maturation, and stress response (Nambara, 2016) and CcpA, which is a core transcriptional regulator in the control of catabolism in Gram-positive bacteria (Carvalho et al., 2011). However, due to the variability in response to MP loading and between replicates (Fig. 6), further research is required to understand the role BAs may play in both quorum sensing and stress regulation in the soil system, as well as their spatial homogeneity.

#### 4.6. Effect of MP on earthworms

Earthworms are key representatives of soil fauna in relation to soil health, performing an important role in the formation and maintenance of soil fertility and structure, as well as being a major contributor to

invertebrate biomass in soil (Blouin et al., 2013). Therefore, understanding the risks that MPs may pose to their health, abundance, and functioning within the agroecosystem is a priority. Earthworms have been shown to transport MPs throughout the soil profile either through adhesion to the exterior of the earthworm body (Rillig et al., 2017b) or egestion of smaller MP particles (Huerta Lwanga et al., 2016). Our study found that there were no significant differences in earthworm abundance or biomass after 6 months of MP incorporation into the soil (Table 1), however, we did not measure egestion or adhesion. This result is inconsistent with much of the existing literature on earthworm exposure to MPs in soil, with several studies reporting negative effects on earthworm physiology (e.g. skin damage, induction of oxidative stress, loss of body weight, reduction in growth, mortality), although experiments were all laboratory or mesocosm based, over short time periods (<60 days) and at maximum loading rates ranging from 1% to 60% (Boots et al., 2019; Cao et al., 2017; Y. H. Chen et al., 2020; Huerta Lwanga et al., 2016; Judy et al., 2019; Rodríguez-Seijo et al., 2019). MP loading rates in the aforementioned experiments ranged from 0.01% to 2% (w/w). Here we added MPs at the rates of 0%, ~0.1%, ~1% and ~10% (w/w), while earthworm health was not directly measured, a lack of change in earthworm abundance or biomass suggests that earthworm health had not diminished significantly, even at high MP loading. By proxy, this also suggests that earthworms do not actively avoid areas of microplastic contamination in the field, as in this study there were no barriers to earthworms leaving the MP loaded plots.

With this, it must be noted that this study only incorporated MPs into the top 10 cm of soil, therefore exposure of earthworms to MPs will likely depend on their ecotype, with endogenic earthworms likely to have higher exposure rates than the deeper dwelling anecic earthworms. As MPs are moved through the soil profile over time it is likely that the full extent of the impact on earthworms will be clearer. Equally, the longer-term (years to decades) impact of MPs is likely to be more severe than the short term. As MP particles degrade and fragment, they will become more ingestible to macrofauna and microfauna, although it is likely that the MP powder added in this study was already small enough to be digestible, possibly leading to greater mortality in soil-dwelling fauna (Lahive et al., 2019). Likewise, earthworms live several years, therefore it is likely that this study captures only a snapshot of the earthworm lifecycle. Longer term monitoring is required to establish trends in earthworm health.

#### 4.7. Crop health as affected by MP loading

The ability to effectively grow healthy crop plants is a key ecosystem service provided by the soil in an agroecosystem context, underpinning human health and nutrition (Power, 2010). However, data on the effect of MP loading on crop yield and health is limited. MPs have the potential to affect plants in several ways; altering the soil structure, immobilising nutrients, contaminant transport, or adsorption and direct toxicity (Rillig et al., 2019). Several short-term laboratory studies have shown the negative effect of MPs on plant health and biomass at loading rates ranging from 0.2 to 2% (de Souza Machado et al., 2019; Y. Qi et al., 2020; Zang et al., 2020). The results of this field study are contradictory to these studies, suggesting that MPs, even at extremely high loading rates, have no significant effects on the aboveground, ear biomass, or C:N ratio of the harvested seed of *T. aestivum* over one cropping season. However, the effect of MPs on root biomass and rooting structure was not measured in this study, though it is likely that the aboveground biomass would be affected if root growth characteristics were altered by MPs, as a high proportion of wheat roots are found within the top 10 cm of soil (Li et al., 2011).

#### 4.8. Implications and future research direction

Most existing data on MPs is based on laboratory or mesocosm based experiments. While these data are useful, field studies better represent

real-world conditions. Longer-term (years to decades) datasets are required to obtain a more comprehensive understanding of the effect of MPs on soil physicochemistry as well as soil biology and plant health. The study of extremely high MP loading rates may also be useful to understand future effects of MP on soil, if continuous loading occurs (e.g. repeated use of plastic mulch films). Generally, it is recommended that loading rates for future MP studies should reflect realistic loading rates in soil to accurately reflect a perturbed system. Even in heavily mulched soil MP loading rarely exceeds 325 kg ha<sup>-1</sup>, although this is likely to increase as MPs continue to be added to the soil (Huang et al., 2020), although little data explicitly reporting loading rates is available, with many studies choosing to report as items kg<sup>-1</sup> (Büks and Kauenjohann, 2020).

It must also be noted that the potential negative impacts of (particularly conventional) MPs on soil and ecosystem health are likely to increase over time as their decomposition rates are extremely slow relative to the rate of entry to the system, leading to a progressive accumulation within soil (Rillig, 2012; Rillig et al., 2017a), potentially becoming persistent organic pollutants. Equally, while biodegradation is possible to a small extent, it is likely MPs relative recalcitrance means that microbes will prefer less energetically expensive C sources, and therefore, biological, co-metabolic, break-down of plastic is unlikely to occur to any great extent in field soils (Ng et al., 2018). That is what our data suggests, i.e. that if there are no additives, once a biofilm has formed on the outside, pure MPs are no different from an inert sand particle. However, this study is also limited in respect the size and shape of MPs applied to the soil, which may not be typical of primary or secondary MPs typically applied to, or found in, soils, which in the case of mulch films are more likely to be thin films or peices as opposed to individual particles applied here (Huang et al., 2020).

This study applied pure MP LDPE powder, with very low levels of contaminants and additives present. The chemical formulation of MP entering agricultural soils, however, is expected to vary widely due to their origin (e.g. mulch film, biosolids) giving rise to variable amounts of additives (co-pollutants) such as plasticisers (generally low-volatility, insoluble and chemically stable; Campanale et al., 2020), colourants and pigments (inorganic pigments containing heavy metals or organic pigments including various chromophoric families that are potentially carcinogenic and mutagenic; Gičević et al., 2020; Völz, 2009), ultraviolet (UV) stabilisers (inorganic or organic cadmium, barium, or lead salts; Stenmarck et al., 2017) or other polymers (Steinmetz et al., 2016). Generally, additives are not chemically bound to the plastic polymer and subsequent leaching of these additives may pose more of a hazard to soil ecology (particularly microorganisms) than the relatively recalcitrant MP themselves, particularly in the short term (days to years). The exchange and effects of additives or contaminants between plastic particles and the surrounding soil environment and the subsequent effect on soil function (e.g. enzyme inhibition) is a key area for future terrestrial plastics research.

It is also important to state that the majority of published literature on MPs does not state the purity of the plastics, MP used and the type (and concentration) of aforementioned additives incorporated. Reporting of this information is highly recommended in future literature, due to the potential varying effects on the soil environment as well as toxicity to soil ecology, which may significantly affect the results, particularly of biological studies.

## 5. Conclusions

This study demonstrated that the application of pure LDPE MP powder to a field site with no previous history of plastic pollution or application had no significant effect on soil biological health or function over one growing season (6 months). In this regard, we reject hypotheses i, ii and iii, as there were no significant changes in biological quality, crop biomass, or yield with MP loading; equally no effect of loading rate was observed. In conclusion, MPs themselves may not pose a significant

problem, at least in the short term (days to years) due to their recalcitrant nature. Further work should be undertaken focusing on the effect of additives and contaminants on soil function and plant health, as well as the longer-term (years to decades) effects of MP incorporation to soil, in a field context.

### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Acknowledgements

We thank Joe Cotton, for his help in the maintenance of the field trial, and Jennifer Rhymes for statistical discussion. This work was initiated using the UKRI Global Challenges Research Fund (GCRF) award made available by the Higher Education Funding Council for Wales (HEFCW) to Bangor University (W19/36HE) and subsequently supported by the GCRF project awarded to Bangor University (NE/V005871/1). We acknowledge use of the Microscopy Australia facilities at UWA, a facility funded by UWA, and State and Federal Governments, and thank Sarah Gain for SEM technical assistance. Robert Brown is supported through a Knowledge Economy Skills Scholarship (KESS 2). KESS 2 is a pan-Wales higher level skills initiative led by Bangor University on behalf of the HE sector in Wales. It is part funded by the Welsh Government's European Social Fund (ESF) convergence programme for West Wales and the Valleys.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.soilbio.2021.108496>.

### References

- AHDB, 2018. Nutrient Management Guide (RB209). Section 4 Arable Crops, p. 52. <https://ahdb.org.uk/knowledge-library/rb209-section-4-arable-crops>.
- Allen, S., Allen, D., Phoenix, V.R., Le Roux, G., Jimenez, P.D., Simonneau, A., Binet, S., Galop, D., 2019. Atmospheric transport and deposition of microplastics in a remote mountain catchment. *Nature Geoscience* 12, 339–344. <https://doi.org/10.1038/s41561-019-0335-5>.
- Ángeles-López, Y.G., Gutiérrez-Mayen, A.M., Velasco-Pérez, M., Beltrán-Villavicencio, M., Vázquez-Morillas, A., Cano-Blanco, M., 2017. Abiotic degradation of plastic films. *Journal of Physics: Conference Series* 792, 012027. <https://doi.org/10.1088/1742-6596/792/1/012027>.
- Avio, C.G., Gorbí, S., Regoli, F., 2017. Plastics and microplastics in the oceans: from emerging pollutants to emerged threat. *Marine Environmental Research* 128, 2–11. <https://doi.org/10.1016/j.marenvres.2016.05.012>.
- Awet, T.T., Kohl, Y., Meier, F., Straskraba, S., Grün, A.L., Ruf, T., Jost, C., Drexel, R., Tunc, E., Emmerling, C., 2018. Effects of polystyrene nanoparticles on the microbiota and functional diversity of enzymes in soil. *Environmental Sciences Europe* 30, 11. <https://doi.org/10.1186/s12302-018-0140-6>.
- Bandopadhyay, S., Sintim, H.Y., DeBruyn, J.M., 2020. Effects of biodegradable plastic film mulching on soil microbial communities in two agroecosystems. *PeerJ* 2020, e9015. <https://doi.org/10.7717/peerj.9015>.
- Barrat, H.A., Evans, J., Chadwick, D.R., Clark, I.M., le Cocq, K., Cardenas, L.M., 2020. The impact of drought and rewetting on N<sub>2</sub>O emissions from soil in temperate and Mediterranean climates. *European Journal of Soil Science*. <https://doi.org/10.1111/ejss.13015>.
- Bates, D., Mächler, M., Bolker, B.M., Walker, S.C., 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67, 1–48. <https://doi.org/10.18637/jss.v067.i01>.
- Bell, M.J., Winning, N., Rees, R.M., Cloy, J.M., Topp, K., Cardenas, L., Donovan, N., Scott, T., Webster, C., Whitmore, A., Williams, J., Balshaw, H., Paine, F., Chadwick, D., 2015. Nitrous Oxide emissions from fertilised UK arable soils: quantification and mitigation. *Agriculture, Ecosystems & Environment* 212, 134–147.
- Blagodatskaya, E., Kuzyakov, Y., 2013. Active microorganisms in soil: critical review of estimation criteria and approaches. *Soil Biology and Biochemistry* 67, 192–211. <https://doi.org/10.1016/j.soilbio.2013.08.024>.
- Bläsing, M., Amelung, W., 2018. Plastics in soil: analytical methods and possible sources. *The Science of the Total Environment* 612, 422–435. <https://doi.org/10.1016/j.scitotenv.2017.08.086>.
- Blouin, M., Hodson, M.E., Delgado, E.A., Baker, G., Brussaard, L., Butt, K.R., Dai, J., Dendooven, L., Peres, G., Tondoh, J.E., Cluzeau, D., Brun, J.-J., 2013. A review of earthworm impact on soil function and ecosystem services. *European Journal of Soil Science* 64, 161–182. <https://doi.org/10.1111/ejss.12025>.
- Boots, B., Russell, C.W., Green, D.S., 2019. Effects of microplastics in soil ecosystems: above and below ground. *Environmental Science and Technology* 53, 11496–11506. <https://doi.org/10.1021/acs.est.9b03304>.
- Bossio, D.A., Scow, K.M., 1998. Impacts of carbon and flooding on soil microbial communities: phospholipid fatty acid profiles and substrate utilization patterns. *Microbial Ecology* 35, 265–278. <https://doi.org/10.1007/s002489900082>.
- Brown, R.W., Bull, I.D., Journeaux, T., Chadwick, D.R., Jones, D.L., 2021. Volatile organic compounds (VOCs) allow sensitive differentiation of biological soil quality. *Soil Biology and Biochemistry* 156, 108187. <https://doi.org/10.1016/j.soilbio.2021.108187>.
- Buchkowski, R.W., Schmitz, O.J., Bradford, M.A., 2015. Microbial stoichiometry overrides biomass as a regulator of soil carbon and nitrogen cycling. *Ecology* 96, 1139–1149. <https://doi.org/10.1890/14-1327.1>.
- Büks, F., Kaupenjohann, M., 2020. Global concentrations of microplastics in soils - a review. *Soil* 6, 649–662. <https://doi.org/10.5194/soil-6-649-2020>.
- Bünemann, E.K., Bongiorno, G., Bai, Z., Creamer, R.E., de Deyn, G., de Goede, R., Flesskens, L., Geissen, V., Kuyper, T.W., Mäder, P., Pulleman, M., Sukkel, W., van Groenigen, J.W., Brussaard, L., 2018. Soil quality – a critical review. *Soil Biology and Biochemistry* 120, 105–125. <https://doi.org/10.1016/j.soilbio.2018.01.030>.
- Buyer, J.S., Sasser, M., 2012. High throughput phospholipid fatty acid analysis of soils. *Applied Soil Ecology* 61, 127–130. <https://doi.org/10.1016/j.apsoil.2012.06.005>.
- Campanale, C., Massarelli, C., Savino, L., Locaputo, V., Uricchio, V.F., 2020. A detailed review study on potential effects of microplastics and additives of concern on human health. *International Journal of Environmental Research and Public Health* 17, 1212. <https://doi.org/10.3390/ijerph17041212>.
- Cao, D., Wang, X., Luo, X., Liu, G., Zheng, H., 2017. Effects of polystyrene microplastics on the fitness of earthworms in an agricultural soil. *IOP Conference Series: Earth and Environmental Science* 61, 012148. <https://doi.org/10.1088/1755-1315/61/1/012148>.
- Cardenas, L.M., Bhogal, A., Chadwick, D.R., McGeough, K., Misselbrook, T., Rees, R.M., Thorman, R.E., Watson, C.J., Williams, J.R., Smith, K.A., Calvet, S., 2019. Nitrogen use efficiency and nitrous oxide emissions from five UK fertilised grasslands. *The Science of the Total Environment* 661, 696–710.
- Carswell, A., Shaw, R., Hunt, J., Sánchez-Rodríguez, A.R., Saunders, K., Cotton, J., Hill, P.W., Chadwick, D.R., Jones, D.L., Misselbrook, T.H., 2018. Assessing the benefits and wider costs of different N fertilisers for grassland agriculture. *Archives of Agronomy and Soil Science* 65, 625–639. <https://doi.org/10.1080/03650340.2018.1519251>.
- Carvalho, S.M., Kloosterman, T.G., Kuipers, O.P., Neves, A.R., 2011. CcpA ensures optimal metabolic fitness of *Streptococcus pneumoniae*. *PLoS One* 6, e26707. <https://doi.org/10.1371/journal.pone.0026707>.
- Chamas, A., Moon, H., Zheng, J., Qiu, Y., Tabassum, T., Jang, J.H., Abu-Omar, M., Scott, S.L., Suh, S., 2020. Degradation rates of plastics in the environment. *ACS Sustainable Chemistry & Engineering* 8, 3494–3511. <https://doi.org/10.1021/acscuschemeng.9b06635>.
- Chang, C., Janzen, H.H., Nakonechny, E.M., Cho, C.M., 1998. Nitrous oxide emission through plants. *Soil Science Society of America Journal* 62, 35–38. <https://doi.org/10.2136/SSSAJ1998.03615995006200010005X>.
- Chen, H., Wang, Y., Sun, X., Peng, Y., Xiao, L., 2020. Mixing effect of poly(lactic acid microplastic) and straw residue on soil property and ecological function. *Chemosphere* 243, 125271. <https://doi.org/10.1016/j.chemosphere.2019.125271>.
- Chen, Y., Liu, X., Leng, Y., Wang, J., 2020. Defense responses in earthworms (*Eisenia fetida*) exposed to low-density polyethylene microplastics in soils. *Ecotoxicology and Environmental Safety* 187, 109788. <https://doi.org/10.1016/j.ecoenv.2019.109788>.
- Chong, J., Soufan, O., Li, C., Caraus, I., Li, S., Bourque, G., Wishart, D.S., Xia, J., 2018. MetaboAnalyst 4.0: towards more transparent and integrative metabolomics analysis. *Nucleic Acids Research* 46, W486–W494. <https://doi.org/10.1093/nar/gky310>.
- Cole, M., Lindeque, P., Halsband, C., Galloway, T.S., 2011. Microplastics as contaminants in the marine environment: a review. *Marine Pollution Bulletin* 62, 2588–2597. <https://doi.org/10.1016/j.marpolbul.2011.09.025>.
- Comerford, N.B., Franzluebbers, A.J., Stromberger, M.E., Morris, L., Markewitz, D., Moore, R., 2013. Assessment and evaluation of soil ecosystem services. *Soil Horizons* 54, 1–14. <https://doi.org/10.2136/sh12-10-0028>.
- Damisch, W., Wiberg, A., 1991. Biomass yield - a topical issue in modern wheat breeding programmes. *Plant Breeding* 107, 11–17. <https://doi.org/10.1111/j.1439-0523.1991.tb00523.x>.
- de Falco, F., di Pace, E., Cocca, M., Avella, M., 2019. The contribution of washing processes of synthetic clothes to microplastic pollution. *Scientific Reports* 9, 1–11. <https://doi.org/10.1038/s41598-019-43023-x>.
- de Souza Machado, A.A., Lau, C.W., Kloas, W., Bergmann, J., Bachelier, J.B., Faltin, E., Becker, R., Görlich, A.S., Rillig, M.C., 2019. Microplastics can change soil properties and affect plant performance. *Environmental Science and Technology* 53, 6044–6052. <https://doi.org/10.1021/acs.est.9b01339>.
- DEFRA, 2019. Farming Statistics Provisional Crop Areas, Yields and Livestock Populations June 2019 - United Kingdom.
- Distaso, M.A., Bargiela, R., Brailsford, F.L., Williams, G.B., Wright, S., Lunev, E.A., Toshchakov, S.V., Yakimov, M.M., Jones, D.L., Golyshin, P.N., Golyshina, O.V., 2020. High representation of archaea across all depths in oxic and low-pH sediment layers underlying an acidic stream. *Frontiers in Microbiology* 11, 576520. <https://doi.org/10.3389/fmicb.2020.576520>.

- Dobbie, K.E., Smith, K.A., 2001. The effects of temperature, water-filled pore space and land use on N<sub>2</sub>O emissions from an imperfectly drained gleysol. *European Journal of Soil Science* 52, 667–673. <https://doi.org/10.1046/j.1365-2389.2001.00395.x>.
- Duncan, D.S., Jewell, K.A., Suen, G., Jackson, R.D., 2016. Detection of short-term cropping system-induced changes to soil bacterial communities differs among four molecular characterization methods. *Soil Biology and Biochemistry* 96, 160–168. <https://doi.org/10.1016/j.soilbio.2016.02.002>.
- Espí, E., Salmerón, A., Fontecha, A., García, Y., Real, A.I., 2006. Plastic films for agricultural applications. *Journal of Plastic Film and Sheeting* 22, 85–102. <https://doi.org/10.1177/8756087906064220>.
- Fadrosch, D.W., Ma, B., Gajer, P., Sengamalay, N., Ott, S., Brotman, R.M., Ravel, J., 2014. An improved dual-indexing approach for multiplexed 16S rRNA gene sequencing on the Illumina MiSeq platform. *Microbiome* 2, 6. <https://doi.org/10.1186/2049-2618-2-6>.
- Fahy, E., Subramaniam, S., 2020. RefMet: a reference nomenclature for metabolomics. *Nature Methods*. <https://doi.org/10.1038/s41592-020-01009-y>.
- Fei, Y., Huang, S., Zhang, H., Tong, Y., Wen, D., Xia, X., Wang, H., Luo, Y., Barceló, D., 2020. Response of soil enzyme activities and bacterial communities to the accumulation of microplastics in an acid cropped soil. *The Science of the Total Environment* 707, 135634. <https://doi.org/10.1016/j.scitotenv.2019.135634>.
- Ferrari, A.E., Ravnskov, S., Larsen, J., Tønnersen, T., Maronna, R.A., Wall, L.G., 2015. Crop rotation and seasonal effects on fatty acid profiles of neutral and phospholipids extracted from no-till agricultural soils. *Soil Use & Management* 31, 165–175. <https://doi.org/10.1111/sum.12165>.
- Fidel, R.B., Laird, D.A., Parkin, T.B., 2019. Effect of biochar on soil greenhouse gas emissions at the laboratory and field scales. *Soil Systems* 3, 8. <https://doi.org/10.3390/soilsystems3010008>.
- Fujihara, S., 2009. Biogenic amines in rhizobia and legume root nodules. *Microbes and Environments* 24, 1–13. <https://doi.org/10.1264/jmsme2.ME08557>.
- Galloway, T.S., Lewis, C.N., 2016. Marine microplastics spell big problems for future generations. *Proceedings of the National Academy of Sciences of the United States of America* 113, 2331–2333. <https://doi.org/10.1073/pnas.1600715113>.
- Gao, H., Yan, C., Liu, Q., Ding, W., Chen, B., Li, Z., 2019. Effects of plastic mulching and plastic residue on agricultural production: a meta-analysis. *The Science of the Total Environment* 651, 484–492. <https://doi.org/10.1016/j.scitotenv.2018.09.105>.
- Gertsman, I., Barshov, B.A., 2018. Promises and pitfalls of untargeted metabolomics. *J. Inher. Metab. Dis.* 41, 355–366. <https://doi.org/10.1007/s10545-017-0130-7>.
- Geyer, R., Jambeck, J.R., Law, K.L., 2017. Production, use, and fate of all plastics ever made. *Sci. Sci. Adv.* 3, e1700782. <https://doi.org/10.1126/sciadv.1700782>.
- Gičević, A., Hindija, L., Karačić, A., 2020. Toxicity of azo dyes in pharmaceutical industry. In: *IFMBE Proceedings*. Springer Verlag, pp. 581–587. [https://doi.org/10.1007/978-3-030-17971-7\\_88](https://doi.org/10.1007/978-3-030-17971-7_88).
- Guo, J.J., Huang, X.P., Xiang, L., Wang, Y.Z., Li, Y.W., Li, H., Cai, Q.Y., Mo, C.H., Wong, M.H., 2020. Source, migration and toxicology of microplastics in soil. *Environment International* 137, 105263. <https://doi.org/10.1016/j.envint.2019.105263>.
- Hou, D., Bolan, N.S., Tsang, D.C.W., Kirkham, M.B., O'Connor, D., 2020. Sustainable soil use and management: an interdisciplinary and systematic approach. *The Science of the Total Environment* 729, 138961. <https://doi.org/10.1016/j.scitotenv.2020.138961>.
- Huang, Y., Liu, Q., Jia, W., Yan, C., Wang, J., 2020. Agricultural plastic mulching as a source of microplastics in the terrestrial environment. *Environmental Pollution* 260, 114096. <https://doi.org/10.1016/j.envpol.2020.114096>.
- Huang, Y., Zhao, Y., Wang, J., Zhang, M., Jia, W., Qin, X., 2019. LDPE microplastic films alter microbial community composition and enzymatic activities in soil. *Environmental Pollution* 254, 112983. <https://doi.org/10.1016/j.envpol.2019.112983>.
- Huerta Lwanga, E., Gertsens, H., Gooren, H., Peters, P., Salánki, T., van der Ploeg, M., Besseling, E., Koelmans, A.A., Geissen, V., 2016. Microplastics in the terrestrial ecosystem: implications for *Lumbricus terrestris* (Oligochaeta, lumbricidae). *Environmental Science and Technology* 50, 2685–2691. <https://doi.org/10.1021/acs.est.5b05478>.
- Hughes, D.T., Sperandio, V., 2008. Inter-kingdom signalling: communication between bacteria and their hosts. *Nature Reviews Microbiology* 6, 111–120. <https://doi.org/10.1038/nrmicro1836>.
- Iqbal, S., Xu, J., Allen, S.D., Khan, S., Nadir, S., Arif, M.S., Yasmeen, T., 2020. Unraveling consequences of soil micro- and nano-plastic pollution on soil-plant system: implications for nitrogen (N) cycling and soil microbial activity. *Chemosphere* 260, 127579. <https://doi.org/10.1016/j.chemosphere.2020.127578>.
- Jia, Y., Whalen, J.K., 2020. A new perspective on functional redundancy and phylogenetic niche conservatism in soil microbial communities. *Pedosphere* 30, 18–24. [https://doi.org/10.1016/S1002-0160\(19\)60826-X](https://doi.org/10.1016/S1002-0160(19)60826-X).
- Jian, J., Xiangbin, Z., Xianbo, H., 2020. An overview on synthesis, properties and applications of poly(butylene-adipate-co-terephthalate)-PBAT. *Adv. Ind. Eng. Polym. Res.* 3, 19–26. <https://doi.org/10.1016/j.aiepr.2020.01.001>.
- Jones, D.L., Rousk, J., Edwards-Jones, G., DeLuca, T.H., Murphy, D.v., 2012. Biochar-mediated changes in soil quality and plant growth in a three year field trial. *Soil Biology and Biochemistry* 45, 113–124. <https://doi.org/10.1016/j.soilbio.2011.10.012>.
- Jost, L., 2010. The relation between evenness and diversity. *Diversity* 2, 207–232. <https://doi.org/10.3390/d2020207>.
- Judy, J.D., Williams, M., Gregg, A., Oliver, D., Kumar, A., Kookana, R., Kirby, J.K., 2019. Microplastics in municipal mixed-waste organic outputs induce minimal short to long-term toxicity in key terrestrial biota. *Environmental Pollution* 252, 522–531. <https://doi.org/10.1016/j.envpol.2019.05.027>.
- Karlen, D.L., Mausbach, M.J., Doran, J.W., Cline, R.G., Harris, R.F., Schuman, G.E., 1997. Soil quality: a concept, definition, and framework for evaluation (A guest editorial). *Soil Science Society of America Journal* 61, 4–10. <https://doi.org/10.2136/sssaj1997.03615995006100010001x>.
- Katsumi, N., Kusube, T., Nagao, S., Okochi, H., 2021. Accumulation of microcapsules derived from coated fertilizer in paddy fields. *Chemosphere* 267, 129185. <https://doi.org/10.1016/j.chemosphere.2020.129185>.
- Kim, D., Chae, Y., An, Y.J., 2017. Mixture toxicity of nickel and microplastics with different functional groups on *Daphnia magna*. *Environmental Science and Technology* 51, 12852–12858. <https://doi.org/10.1021/acs.est.7b03732>.
- Kopittke, P.M., Menzies, N.W., Wang, P., McKenna, B.A., Lombi, E., 2019. Soil and the intensification of agriculture for global food security. *Environment International* 132, 105078. <https://doi.org/10.1016/j.envint.2019.105078>.
- Kuyper, M.M.M., Marchant, H.K., Kartal, B., 2018. The microbial nitrogen-cycling network. *Nature Reviews Microbiology* 16, 263–276. <https://doi.org/10.1038/nrmicro.2018.9>.
- Lahive, E., Walton, A., Horton, A.A., Spurgeon, D.J., Svendsen, C., 2019. Microplastic particles reduce reproduction in the terrestrial worm *Enchytraeus crypticus* in a soil exposure. *Environmental Pollution* 255, 113174. <https://doi.org/10.1016/j.envpol.2019.113174>.
- Lal, R., 2016. Soil health and carbon management. *Food and Energy Secur.* 5, 212–222. <https://doi.org/10.1002/fes3.96>.
- Lau, J.A., Lennon, J.T., 2012. Rapid responses of soil microorganisms improve plant fitness in novel environments. *Proceedings of the National Academy of Sciences of the United States of America* 109, 14058–14062. <https://doi.org/10.1073/pnas.1202319109>.
- Lehman, R.M., Cambardella, C., Stott, D., Acosta-Martinez, V., Manter, D., Buyer, J., Maul, J., Smith, J., Collins, H., Halvorson, J., Kremer, R., Lundgren, J., Ducey, T., Jin, V., Karlen, D., 2015a. Understanding and enhancing soil biological health: the solution for reversing soil degradation. *Sustainability* 7, 988–1027. <https://doi.org/10.3390/su7010988>.
- Lehman, R.M., Acosta-Martinez, V., Buyer, J.S., Cambardella, C.A., Collins, H.P., Ducey, T.F., Halvorson, J.J., Jin, V.L., Johnson, J.M.F., Kremer, R.J., Lundgren, J.G., Manter, D.K., Maul, J.E., Smith, J.L., Stott, D.E., 2015b. Soil biology for resilient, healthy soil. *Journal of Soil and Water Conservation* 70, 12A–18A. <https://doi.org/10.2489/jswc.70.1.12A>.
- Lenth, R.V., 2021. Emmeans: estimated marginal means, aka least-squares means. R package version 1.5.5-1. <https://CRAN.R-project.org/package=emmeans>.
- Li, D., Liu, Y., Chen, Y., Tang, L., Tan, F., Jiang, H., Lei, X., Cao, W., Zhu, Y., 2011. Root architecture modeling and visualization in wheat. *IFIP AICT* 345, 479–490.
- Lin, D., Yang, G., Dou, P., Qian, S., Zhao, L., Yang, Y., Fanin, N., 2020. Microplastics negatively affect soil fauna but stimulate microbial activity: insights from a field-based microplastic addition experiment. *Proceedings of the Royal Society B: Biological Sciences* 287, 20201268. <https://doi.org/10.1098/rspb.2020.1268>.
- Mah, J.H., Park, Y.K., Jin, Y.H., Lee, J.H., Hwang, H.J., 2019. Bacterial production and control of biogenic amines in Asian fermented soybean foods. *Foods* 8, 85. <https://doi.org/10.3390/foods8020085>.
- Marsden, K., Holmberg, J., Jones, D.L., Chadwick, D., 2018. Sheep urine patch N<sub>2</sub>O emissions are lower from extensively-managed than intensively-managed grasslands. *Agriculture, Ecosystems Environ.* 265, 264–274. <https://doi.org/10.1016/j.agee.2018.06.025>.
- McCormick, A., Hoellein, T.J., Mason, S.A., Schluep, J., Kelly, J.J., 2014. Microplastic is an abundant and distinct microbial habitat in an urban river. *Environmental Science and Technology* 48, 11863–11871. <https://doi.org/10.1021/es503610r>.
- Miranda, K.M., Espey, M.G., Wink, D.A., 2001. Spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 5, 62–71. <https://doi.org/10.1006/niox.2000.0319>.
- Moore-Kucera, J., Dick, R.P., 2008. PLFA Profiling of microbial community structure and seasonal shifts in soils of a Douglas-fir chronosequence. *Microbial Ecology* 55, 500–511. <https://doi.org/10.1007/s00428-007-9295-1>.
- Mulvaney, R.L., 1996. Nitrogen - inorganic forms. In: Sparks, D.L. (Ed.), *Methods of Soil Analysis, Part 3*. Soil Science Society of America, Madison, WI, USA, pp. 1123–1184.
- Nambara, E., 2016. Abscisic acid. In: *Encyclopedia of Applied Plant Sciences*. Elsevier Inc., pp. 361–366. <https://doi.org/10.1016/B978-0-12-394807-6.00098-8>.
- Ng, E.L., Huerta Lwanga, E., Eldridge, S.M., Johnston, P., Hu, H.W., Geissen, V., Chen, D., 2018. An overview of microplastic and nanoplastics pollution in agroecosystems. *The Science of the Total Environment* 627, 1377–1388. <https://doi.org/10.1016/j.scitotenv.2018.01.341>.
- Nizzetto, L., Futter, M., Langaas, S., 2016a. Are agricultural soils dumps for microplastics of urban origin? *Environmental Science and Technology* 50, 10777–10779. <https://doi.org/10.1021/acs.est.6b04140>.
- Nizzetto, L., Langaas, S., Futter, M., 2016b. Pollution: do microplastics spill on to farm soils? *Nature* 537, 488. <https://doi.org/10.1038/537488b>.
- North, J.A., Miller, A.R., Wildenthal, J.A., Young, S.J., Robert Tabita, F., 2017. Microbial pathway for anaerobic 5'-methylthioadenosine metabolism coupled to ethylene formation. *Proceedings of the National Academy of Sciences of the United States of America* 114, E10455–E10464. <https://doi.org/10.1073/pnas.1711625114>.
- Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2020. *Vegan: Community Ecology Package*. R Package Version 2.5-7.
- Or, D., Smets, B.F., Wraith, J.M., Dechesne, A., Friedmann, S.P., 2006. Physical constraints affecting bacterial habitats and activity in unsaturated porous media—a review. *Advances in Water Resources* 30, 1505–1527. <https://doi.org/10.1016/j.advwatres.2006.05.025>.
- Ouyang, Y., Evans, S.E., Friesen, M.L., Tiemann, L.K., 2018. Effect of nitrogen fertilization on the abundance of nitrogen cycling genes in agricultural soils: a meta-

- analysis of field studies. *Soil Biology and Biochemistry* 127, 71–78. <https://doi.org/10.1016/j.soilbio.2018.08.024>.
- Pang, Z., Chong, J., Li, S., Xia, J., 2020. MetaboAnalystR 3.0: toward an optimized workflow for global metabolomics. *Metabolites* 10, 186. <https://doi.org/10.3390/metabo10050186>.
- Pérez-Álvarez, E.P., Garde-Cerdán, T., Cabrita, M.J., García-Escudero, E., Peregrina, F., 2017. Influence on wine biogenic amine composition of modifications to soil N availability and grapevine N by cover crops. *Journal of the Science of Food and Agriculture* 97, 4800–4806. <https://doi.org/10.1002/jsfa.8349>.
- Piehl, S., Leibner, A., Löder, M.G.J., Dris, R., Bogner, C., Laforsch, C., 2018. Identification and quantification of macro- and microplastics on an agricultural farmland. *Scientific Reports* 8, 17950. <https://doi.org/10.1038/s41598-018-36172-y>.
- Power, A.G., 2010. Ecosystem services and agriculture: tradeoffs and synergies. *Philosophical Transactions of the Royal Society of London - Series B: Biological Sciences* 365, 2959–2971. <https://doi.org/10.1098/rstb.2010.0143>.
- Qi, R.M., Jones, D.L., Li, Z., Liu, Q., Yan, C.R., 2020. Behavior of microplastics and plastic film residues in the soil environment: a critical review. *The Science of the Total Environment* 703, 134722. <https://doi.org/10.1016/j.scitotenv.2019.134722>.
- Qi, Y., Ossowicki, A., Yang, X., Huerta Lwanga, E., Dini-Andreote, F., Geissen, V., Garbeva, P., 2020. Effects of plastic mulch film residues on wheat rhizosphere and soil properties. *Journal of Hazardous Materials* 387, 121711. <https://doi.org/10.1016/j.jhazmat.2019.121711>.
- Quilliam, R.S., Glanville, H.C., Wade, S.C., Jones, D.L., 2013. Life in the “charosphere” - does biochar in agricultural soil provide a significant habitat for microorganisms? *Soil Biology and Biochemistry* 65, 287–293. <https://doi.org/10.1016/j.soilbio.2013.06.004>.
- R Core Team, 2021. *R: A Language and Environment for Statistical Computing*.
- Ren, X., Tang, J., Liu, X., Liu, Q., 2020. Effects of microplastics on greenhouse gas emissions and the microbial community in fertilized soil. *Environmental Pollution* 256, 113347. <https://doi.org/10.1016/j.envpol.2019.113347>.
- Rillig, M.C., 2012. Microplastic in terrestrial ecosystems and the soil? *Environmental Science and Technology*. <https://doi.org/10.1021/es302011r>.
- Rillig, M.C., Hoffmann, M., Lehmann, A., Liang, Y., Lück, M., Augustin, J., 2021. Microplastic fibers affect dynamics and intensity of CO<sub>2</sub> and N<sub>2</sub>O fluxes from soil differently. *Microplastics and Nanoplastics* 1 (1 1), 1–11. <https://doi.org/10.1186/S43591-021-00004-0>, 2021.
- Rillig, M.C., Ingrassia, R., de Souza Machado, A.A., 2017a. Microplastic incorporation into soil in agroecosystems. *Frontiers of Plant Science* 8, 1805. <https://doi.org/10.3389/fpls.2017.01805>.
- Rillig, M.C., Lehmann, A., Souza Machado, A.A., Yang, G., 2019. Microplastic effects on plants. *New Phytologist* 223, 1066–1070. <https://doi.org/10.1111/nph.15794>.
- Rillig, M.C., Ziersch, L., Hempel, S., 2017b. Microplastic transport in soil by earthworms. *Scientific Reports* 7, 1–6. <https://doi.org/10.1038/s41598-017-01594-7>.
- Rinot, O., Levy, G.J., Steinberger, Y., Svoray, T., Eshel, G., 2019. Soil health assessment: a critical review of current methodologies and a proposed new approach. *The Science of the Total Environment* 648, 1484–1491. <https://doi.org/10.1016/j.scitotenv.2018.08.259>.
- Rocha-Santos, T., Duarte, A.C., 2015. A critical overview of the analytical approaches to the occurrence, the fate and the behavior of microplastics in the environment. *TRAC Trends in Analytical Chemistry* 65, 47–53. <https://doi.org/10.1016/j.trac.2014.10.011>.
- Rodríguez-Sejido, A., Santos, B., Ferreira da Silva, E., Cachada, A., Pereira, R., 2019. Low-density polyethylene microplastics as a source and carriers of agrochemicals to soil and earthworms. *Environmental Chemistry* 16, 8–17. <https://doi.org/10.1071/EN18162>.
- Rong, L., Zhao, Longfei, Zhao, Leicheng, Cheng, Z., Yao, Y., Yuan, C., Wang, L., Sun, H., 2021. LDPE microplastics affect soil microbial communities and nitrogen cycling. *The Science of the Total Environment* 773, 145640. <https://doi.org/10.1016/j.scitotenv.2021.145640>.
- Ruiz-Capillas, C., Herrero, A., 2019. Impact of biogenic amines on food quality and safety. *Foods* 8, 62. <https://doi.org/10.3390/foods8020062>.
- Scheer, C., Rowlings, D.W., Firrel, M., Deuter, P., Morris, S., Grace, P.R., 2014. Impact of nitrification inhibitor (DMPP) on soil nitrous oxide emissions from an intensive broccoli production system in sub-tropical Australia. *Soil Biology and Biochemistry* 77, 243–251. <https://doi.org/10.1016/j.soilbio.2014.07.006>.
- Schimel, J.P., 2018. Life in dry soils: effects of drought on soil microbial communities and processes. *Annual Review of Ecology Evolution and Systematics* 12, 409–432. <https://doi.org/10.1146/annurev-ecolsys-110617>.
- Seeley, M.E., Song, B., Passie, R., Hale, R.C., 2020. Microplastics affect sedimentary microbial communities and nitrogen cycling. *Nature Communications* 11, 1–10. <https://doi.org/10.1038/s41467-020-16235-3>.
- Sharma, S., Chatterjee, S., 2017. Microplastic pollution, a threat to marine ecosystem and human health: a short review. *Environmental Science and Pollution Research* 24, 21530–21547. <https://doi.org/10.1007/s11356-017-9910-8>.
- Sebille, E. van, Wilcox, C., Lebreton, L., Maximenko, N., Hardesty, B.D., Franeker, J.A. van, Eriksen, M., Siegel, D., Galgani, F., Law, K.L., 2015. A global inventory of small floating plastic debris. *Environmental Research Letters* 10, 124006. <https://doi.org/10.1088/1748-9326/10/12/124006>.
- Shen, M., Huang, W., Chen, M., Song, B., Zeng, G., Zhang, Y., 2020. (Micro)plastic crisis: un-ignorable contribution to global greenhouse gas emissions and climate change. *Journal of Cleaner Production* 254, 120138. <https://doi.org/10.1016/j.jclepro.2020.120138>.
- Steinmetz, Z., Wollmann, C., Schaefer, M., Buchmann, C., David, J., Tröger, J., Muñoz, K., Frör, O., Schaumann, G.E., 2016. Plastic mulching in agriculture. Trading short-term agronomic benefits for long-term soil degradation? *The Science of the Total Environment* 550, 690–705. <https://doi.org/10.1016/j.scitotenv.2016.01.153>.
- Stenmarck, Å., Belleza, E.L., Fråne, A., Busch, N., Larsen, Å., Wahlström, M., 2017. Hazardous Substances in Plastics. Nordic Council of Ministers, Copenhagen. <https://doi.org/10.6027/TN2017-505>.
- Stocker, T., 2014. *Climate Change 2013: the Physical Science Basis. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, New York.
- Sudo, N., 2019. Biogenic amines: signals between commensal microbiota and gut physiology. *Frontiers in Endocrinology* 10, 504. <https://doi.org/10.3389/fendo.2019.00504>.
- Sun, Y., Ren, X., Pan, J., Zhang, Z., Tsui, T.H., Luo, L., Wang, Q., 2020. Effect of microplastics on greenhouse gas and ammonia emissions during aerobic composting. *The Science of the Total Environment* 737, 139856. <https://doi.org/10.1016/j.scitotenv.2020.139856>.
- Tosi, M., Brown, S., Ferrari Machado, P.V., Wagner-Riddle, C., Dunfield, K., 2020. Short-term response of soil N-cycling genes and transcripts to fertilization with nitrification and urease inhibitors, and relationship with field-scale N<sub>2</sub>O emissions. *Soil Biology and Biochemistry* 142, 107703. <https://doi.org/10.1016/j.soilbio.2019.107703>.
- Tsugawa, H., Cajka, T., Kind, T., Ma, Y., Higgins, B., Ikeda, K., Kanazawa, M., Vanderghenst, J., Fiehn, O., Arita, M., 2015. MS-DIAL: data-independent MS/MS deconvolution for comprehensive metabolome analysis. *Nature Methods* 12, 523–526. <https://doi.org/10.1038/nmeth.3393>.
- van den Berg, P., Huerta-Lwanga, E., Corradini, F., Geissen, V., 2020. Sewage sludge application as a vehicle for microplastics in eastern Spanish agricultural soils. *Environmental Pollution* 261, 114198. <https://doi.org/10.1016/j.envpol.2020.114198>.
- Vitousek, P.M., Howarth, R.W., 1991. Nitrogen limitation on land and in the sea: how can it occur? *Biogeochemistry* 13, 87–115. <https://doi.org/10.1007/BF00002772>.
- Völz, H.G., 2009. Pigments, inorganic, 1. General. In: *Ullmann's Encyclopedia of Industrial Chemistry*. Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany. [https://doi.org/10.1002/14356007.a20\\_243.pub3](https://doi.org/10.1002/14356007.a20_243.pub3).
- Watteau, F., Dignac, M.-F., Bouchard, A., Revallier, A., Houot, S., 2018. Microplastic Detection in soil amended with municipal solid waste composts as revealed by transmission electronic microscopy and pyrolysis/GC/MS. *Frontiers in Sustainable Food Syst.* 2, 81. <https://doi.org/10.3389/fsufs.2018.00081>.
- Wickham, H., 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New York.
- Withers, E., Hill, P.W., Chadwick, D.R., Jones, D.L., 2020. Use of untargeted metabolomics for assessing soil quality and microbial function. *Soil Biology and Biochemistry* 143, 107758. <https://doi.org/10.1016/j.soilbio.2020.107758>.
- Yan, Y., Chen, Z., Zhu, F., Zhu, C., Wang, C., Gu, C., 2020. Effect of polyvinyl chloride microplastics on bacterial community and nutrient status in two agricultural soils. *Bulletin of Environmental Contamination and Toxicology* 1, 3. <https://doi.org/10.1007/s00128-020-02900-2>.
- Yi, M., Zhou, S., Zhang, L., Ding, S., 2021. The effects of three different microplastics on enzyme activities and microbial communities in soil. *Water Environment Research* 93, 24–32. <https://doi.org/10.1002/wer.1327>.
- Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W., Glöckner, F.O., 2014. The SILVA and “all-species living tree project (LTP)” taxonomic frameworks. *Nucleic Acids Research* 42, D643. <https://doi.org/10.1093/nar/gkt1209>.
- Zang, H., Zhou, J., Marshall, M.R., Chadwick, D.R., Wen, Y., Jones, D.L., 2020. Microplastics in the agroecosystem: are they an emerging threat to the plant-soil system? *Soil Biology and Biochemistry* 148, 107926. <https://doi.org/10.1016/j.soilbio.2020.107926>.
- Zhang, M., Zhao, Y., Qin, X., Jia, W., Chai, L., Huang, M., Huang, Y., 2019. Microplastics from mulching film is a distinct habitat for bacteria in farmland soil. *The Science of the Total Environment* 688, 470–478. <https://doi.org/10.1016/j.scitotenv.2019.06.108>.